

**Table 14. HIV infection cases<sup>1</sup> in adolescents and adults under age 25, by sex and exposure category, reported through December 1998, from the 30 areas with confidential HIV infection reporting<sup>2</sup>**

Male exposure category	13-19 years old				20-24 years old			
	1998		Cumulative total		1998		Cumulative total	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Men who have sex with men	145	(52)	911	(48)	575	(50)	5,475	(55)
Injecting drug use	7	(3)	99	(5)	50	(4)	562	(6)
Men who have sex with men and inject drugs	4	(1)	95	(5)	53	(5)	655	(7)
Hemophilia/coagulation disorder	5	(2)	104	(5)	4	(0)	83	(1)
Heterosexual contact:	19	(7)	134	(7)	88	(8)	614	(6)
<i>Sex with injecting drug user</i>	2		25		9		94	
<i>Sex with person with hemophilia</i>	—		2		—		1	
<i>Sex with transfusion recipient with HIV infection</i>	—		1		1		7	
<i>Sex with HIV-infected person, risk not specified</i>	17		106		78		512	
Receipt of blood transfusion, blood components, or tissue	—	—	9	(0)	1	(0)	30	(0)
Risk not reported or identified <sup>3</sup>	99	(35)	542	(29)	368	(32)	2,454	(25)
Male subtotal	279	(100)	1,894	(100)	1,139	(100)	9,873	(100)
<b>Female exposure category</b>								
Injecting drug use	16	(4)	158	(7)	42	(5)	611	(13)
Hemophilia/coagulation disorder	—	—	—	—	1	(0)	3	(0)
Heterosexual contact:	183	(41)	1,106	(49)	313	(36)	2,117	(45)
<i>Sex with injecting drug user</i>	21		221		43		545	
<i>Sex with bisexual male</i>	7		73		16		192	
<i>Sex with person with hemophilia</i>	3		19		—		30	
<i>Sex with transfusion recipient with HIV infection</i>	—		4		2		17	
<i>Sex with HIV-infected person, risk not specified</i>	152		789		252		1,333	
Receipt of blood transfusion, blood components, or tissue	—	—	15	(1)	2	(0)	26	(1)
Risk not reported or identified	249	(56)	986	(44)	502	(58)	1,998	(42)
Female subtotal	448	(100)	2,265	(100)	860	(100)	4,755	(100)
<b>Total<sup>4</sup></b>	<b>728</b>		<b>4,161</b>		<b>1,999</b>		<b>14,628</b>	

<sup>1</sup>Includes only persons reported with HIV infection who have not developed AIDS.

<sup>2</sup>See table 3 for areas with confidential HIV infection reporting for adults and adolescents.

<sup>3</sup>For HIV infection cases, "risk not reported or identified" refers primarily to persons whose mode of exposure was not reported and who have not been followed up to determine their mode of exposure, and to a smaller number of persons who are not reported with one of the exposures listed above after follow-up. See Technical Notes.

<sup>4</sup>Includes 2 persons whose sex is unknown.